

Clinical Policy: Posterior Tibial Nerve Stimulation for Voiding Dysfunction

Reference Number: LA.CP.MP.133

Date of Last Revision: 08/24

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Posterior tibial nerve stimulation (PTNS), also known as peripheral tibial nerve stimulation, is a minimally invasive form of electrical neuromodulation used to treat overactive bladder (OAB) syndrome and associated symptoms of urinary urgency, urinary frequency, and urge urinary incontinence.¹ This policy describes the medical necessity requirements for posterior tibial nerve stimulation.

Policy/Criteria

- I. It is the policy of Louisiana Healthcare Connections that posterior tibial nerve stimulation (PTNS) is medically necessary when all of the following criteria are met:
 - A. Diagnosis of overactive bladder (OAB);
 - B. There has been a failure of conservative medical management (e.g. behavioral therapies such as bladder training or pelvic floor muscle training, or pharmacotherapy with oral medications used with the intent to treat OAB), unless conservative management is not desired or is medically contraindicated;
 - C. Service is provided in accordance with the standard treatment regimen of 30-minute weekly sessions for 12 weeks.
- II. It is the policy of Louisiana Healthcare Connections that once-a-month maintenance treatments with PTNS **are medically necessary** for patients who experience significant improvement in their OAB symptoms after the 12 initial treatments. Treatment frequency may vary depending on return of symptoms.
- III. It is the policy of Louisiana Healthcare Connections that there is insufficient evidence to support the use of PTNS beyond 12 months or when there is no improvement in urinary dysfunction.
- IV. It is the policy of Louisiana Healthcare Connections that there is insufficient evidence in the published peer-reviewed literature to support the use of implantable tibial nerve stimulation for the treatment of urinary voiding dysfunction.

Background

The term "voiding dysfunction" has been used to refer to urinary incontinence, urinary retention, and symptoms of frequency and urgency. Overactive bladder (OAB) is a specific type of voiding dysfunction that includes any of the following symptoms: urinary frequency, urinary urgency, urge incontinence, and nocturia.² OAB can significantly impact quality of life including physical function, sexual function, and social interactions. Treatments for OAB include lifestyle changes, bladder training, pelvic floor muscle training and anticholinergic (anti-muscarinic) drugs.³

Posterior tibial nerve stimulation (PTNS) involves indirect modulation of the specific nerve that controls bladder function (i.e., the sacral nerve plexus) via stimulation of the posterior tibial nerve accessed just above the ankle. This minimally invasive form of neuromodulation consists of insertion of a 34-gauge needle electrode approximately five centimeters (cm) cephalad to the medial malleolus and two cm posterior to the tibia near the tibial nerve. A surface electrode is placed on the medial aspect of the foot. The needle electrode is connected via a lead wire to a low-voltage electrical stimulator. Stimulation is administered at a current level of 0.5 to nine milliamperes (mA) at 20 hertz (Hz) and continues for 30 minutes. Initial treatment regimens typically consist of 12 weekly sessions, with responders exhibiting some symptom improvement after six to eight sessions. Maintenance treatment sessions may be required to sustain the response to treatment.⁴

Several implantable tibial nerve neuromodulation systems, including a battery-less leadless, miniature implantable device, are currently under investigation for the management of OAB, however, evidence is still limited on their benefits and efficacy at this time.

National Institute for Health and Care Excellence (NICE)

According to NICE, current evidence demonstrates that PTNS for OAB syndrome is effective in reducing symptoms in the short term and medium term. Per NICE guidance, PTNS for OAB syndrome does not have major safety concerns, and the use of this procedure should with standard protocols for consent, audit, and clinical governance.³

A NICE guidance on urinary incontinence in women does not recommend the “routine” use of PTNS to treat OAB. Rather, they recommend PTNS for OAB for following:

- There has been a multidisciplinary team (MDT) review, and
- Conservative management including OAB drug treatment has not worked adequately, and
- The woman does not want botulinum toxin A or percutaneous sacral nerve stimulation.¹⁰

American Urological Association

Clinicians may offer PTNS as third-line treatment in a carefully selected patient population, characterized by moderately severe baseline incontinence and frequency and willingness to comply with the PTNS protocol. Patients must also have the resources to make frequent office visits both during the initial treatment phase and to obtain maintenance treatments in order to achieve and maintain treatment effects. The most common protocol is the application of 30 minutes of stimulation once a week for 12 weeks (the trial duration; for continued benefit, weekly stimulation would have to continue).¹

Studies to date evaluating PTNS for the treatment of OAB conclude there is evidence of benefit, although most studies have been small and report short-term outcomes after 12 weeks of treatment. A small study of 33 PTNS responders who continued therapy for six to 12 months reported excellent durability through 12 months.⁵ Another small study reported sustained safety and efficacy of PTNS for the treatment of OAB symptom control over 24 months with initial success after 12 weekly treatments, followed by a 14-week prescribed tapering protocol and a personalized treatment plan with an average of 1.3 treatments per month.⁶

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2023, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

NOTE: Coverage is subject to each requested code’s inclusion on the corresponding LDH fee schedule. Non-covered codes are denoted (*) and are reviewed for Medical Necessity for members under 21 years of age on a per case basis.

CPT codes that support medical necessity

CPT® Codes	Description
64566	Posterior tibial neurostimulation, percutaneous needle electrode, single treatment, includes programming

CPT codes that do not support medical necessity

CPT® Codes	Description
0587T*	Percutaneous implantation or replacement of integrated single device neurostimulation system including electrode array and receiver or pulse generator, including analysis, programming and imaging guidance when performed, posterior tibial nerve
0588T*	Revision or removal of integrated single device neurostimulation system including electrode array and receiver or pulse generator, including analysis, programming, and imaging guidance when performed, posterior tibial nerve
0589T*	Electronic analysis with simple programming of implanted integrated neurostimulation system (eg, electrode array and receiver), including contact group(s), amplitude, pulse width, frequency (Hz), on/off cycling, burst, dose lockout, patient-selectable parameters, responsive neurostimulation, detection algorithms, closed-loop parameters, and passive parameters, when performed by physician or other qualified health care professional, posterior tibial nerve, 1-3 parameters
0590T*	Electronic analysis with complex programming of implanted integrated neurostimulation system (eg, electrode array and receiver), including contact group(s), amplitude, pulse width, frequency (Hz), on/off cycling, burst, dose lockout, patient-selectable parameters, responsive neurostimulation, detection algorithms, closed-loop parameters, and passive parameters, when performed by physician or other qualified health care professional, posterior tibial nerve, 4 or more parameters

Reviews, Revisions, and Approvals	Revision Date	Approval Date	Effective Date
Converted corporate to local policy.	10/20		
Annual review. Replaced “investigational” language with “insufficient evidence to support.” References reviewed, reformatted, and updated. Changed “review date” in the header to “date of last revision” and “date” in the revision log header to “revision date.” Replaced member with member/enrollee. Added “may not support medical necessity” to coding implications. Specialist review.	2/22	2/22	
Annual review. Revised Criteria I.B. to include examples of behavioral therapies such as bladder training or pelvic floor muscle training. Background updated to with no impact on criteria. Dashes removed from code ranges. References reviewed and updated.	9/22		
Annual review. Revised policy statement and all criteria verbiage in criteria I. ICD-10 CM Diagnosis Code table removed. References reviewed and updated. Reviewed by external specialist. Note for non-covered codes added.	09/23	11/27/23	
Annual review. Updated criteria under I.B. by replacing anti-muscarinics or β 3-adrenoceptor agonists and/or antibiotics for urinary tract infections with medications used with the intent to treat OAB. References reviewed and updated.	08/24	10/23/24	11/22/24

References

1. Lightner DJ, Gomelsky A, Souter L, Vasavada SP. Diagnosis and Treatment of Overactive Bladder (Non-Neurogenic) in Adults: AUA/SUFU Guideline Amendment 2019. *J Urol.* 2019;202(3):558 to 563. doi:10.1097/JU.0000000000000309
2. Lukacz ES. Urgency urinary incontinence/overactive bladder (OAB) in females: Treatment. UpToDate. www.uptodate.com. Published July 11, 2023. Accessed June 6, 2024.
3. National Institute for Health and Care Excellence. Percutaneous posterior tibial nerve stimulation for overactive bladder syndrome - Interventional procedures guidance [IPG362]. <https://www.nice.org.uk/guidance/ipg362>. Published October 2010. Accessed June 6, 2024.
4. Health Technology Assessment. Comparative effectiveness review of percutaneous tibial nerve stimulation for the treatment of symptomatic non-neurogenic overactive bladder. Hayes. www.hayesinc.com. Published October 31, 2018 (annual review November 15, 2022) Accessed June 6, 2024.
5. MacDiarmid SA, Peters KM, Shobeiri SA, et al. Long-term durability of percutaneous tibial nerve stimulation for the treatment of overactive bladder. *J Urol.* 2010;183(1):234 to 240. doi:10.1016/j.juro.2009.08.160
6. Peters KM, Carrico DJ, MacDiarmid SA, et al. Sustained therapeutic effects of percutaneous tibial nerve stimulation: 24 month results of the STEP Study. *Neurourol Urodyn.* 2013;32(1):24 to 29. doi:10.1002/nau.22266
7. Peters KM, Macdiarmid SA, Wooldridge LS, et al. Randomized trial of percutaneous tibial nerve stimulation versus extended-release tolterodine: results from the overactive bladder innovative therapy trial. *J Urol.* 2009;182(3):1055 to 1061. doi:10.1016/j.juro.2009.05.045

8. Ammi M, Chautard D, Brassart E, Culty T, Azzouzi AR, Bigot P. Transcutaneous posterior tibial nerve stimulation: evaluation of a therapeutic option in the management of anticholinergic refractory overactive bladder. *Int Urogynecol J*. 2014;25(8):1065 to 1069. doi:10.1007/s00192-014-2359-0
9. Yoong W, Shah P, Dadswell R, Green L. Sustained effectiveness of percutaneous tibial nerve stimulation for overactive bladder syndrome: 2 year follow up of positive responders. *Int Urogynecol J*. 2013;24(5):795 to 799. doi:10.1007/s00192-012-1936-3
10. National Institute for Health and Care Excellence. Urinary incontinence and pelvic organ prolapse in women: management - NICE guideline [NG123]. <https://www.nice.org.uk/guidance/ng123>. Published April 02, 2019. (Last updated June 24, 2019). Accessed June 6, 2024.
11. Johnson TM. Nocturia: Clinical presentation, evaluation, and management in adults. UpToDate. www.uptodate.com. Published February 15, 2023. Accessed June 6, 2024.
12. Blue Cross Blue Shield; Kaiser Foundation Health Plan; Southern California Permanente Medical Group. Percutaneous tibial nerve stimulation for the treatment of voiding dysfunction. *Technol Eval Cent Assess Program Exec Summ*. 2014;28(10):1 to 12.
13. van Breda HMK, Martens FMJ, Tromp J, Heesakkers JPFA. A New Implanted Posterior Tibial Nerve Stimulator for the Treatment of Overactive Bladder Syndrome: 3-Month Results of a Novel Therapy at a Single Center. *J Urol*. 2017;198(1):205 to 210. doi:10.1016/j.juro.2017.01.078
14. Del Río-Gonzalez S, Aragon IM, Castillo E, et al. Percutaneous Tibial Nerve Stimulation Therapy for Overactive Bladder Syndrome: Clinical Effectiveness, Urodynamic, and Durability Evaluation. *Urology*. 2017;108:52 to 58. doi:10.1016/j.urology.2017.04.059
15. Health Technology Assessment. Percutaneous tibial nerve stimulation for the treatment of symptomatic neurogenic lower urinary tract dysfunction. Hayes. www.hayesinc.com. Published April 15, 2019 (annual review April 01, 2022) Accessed June 6, 2024.
16. Gaziev G, Topazio L, Iacovelli V, et al. Percutaneous Tibial Nerve Stimulation (PTNS) efficacy in the treatment of lower urinary tract dysfunctions: a systematic review. *BMC Urol*. 2013;13:61. Published 2013 Nov 25. doi:10.1186/1471-2490-13-61
17. Yamashiro J, de Riese W, de Riese C. New Implantable Tibial Nerve Stimulation Devices: Review of Published Clinical Results in Comparison to Established Neuromodulation Devices. *Res Rep Urol*. 2019;11:351 to 357. Published 2019 Dec 23. doi:10.2147/RRU.S231954
18. Vecchioli-Scaldazza C, Morosetti C, Berouz A, Giannubilo W, Ferrara V. Solifenacin succinate versus percutaneous tibial nerve stimulation in women with overactive bladder syndrome: results of a randomized controlled crossover study. *Gynecol Obstet Invest*. 2013;75(4):230 to 234. doi:10.1159/000350216
19. van der Pal F, van Balken MR, Heesakkers JP, Debruyne FM, Kiemeneij LA, Bemelmans BL. Correlation between quality of life and voiding variables in patients treated with percutaneous tibial nerve stimulation. *BJU Int*. 2006;97(1):113 to 116. doi:10.1111/j.1464-410X.2006.05860.x

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program

approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of member/enrollees. This clinical policy is not intended to recommend treatment for member/enrollees. Member/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, member/enrollees and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, member/enrollees and their representatives agree to be bound by such terms and conditions by providing services to member/enrollees and/or submitting claims for payment for such services.

©2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare

CLINICAL POLICY

Posterior Tibial Nerve Stimulation for Voiding Dysfunction



Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.