

Clinical Policy: Ramucirumab (Cyramza)

Reference Number: LA.PHAR.119

Effective Date: 04.28.21

Last Review Date: 05.02.24

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

****Please note: This policy is for medical benefit****

Description

Ramucirumab (Cyramza®) is a human vascular endothelial growth factor receptor 2 (VEGFR2) antagonist.

FDA Approved Indication(s)

Cyramza is indicated:

- As a single agent or in combination with paclitaxel, for treatment of advanced or metastatic gastric or gastro-esophageal junction (i.e., esophagogastric junction; EGJ) adenocarcinoma, with disease progression on or after prior fluoropyrimidine- or platinum-containing chemotherapy.
- In combination with erlotinib, for first-line treatment of metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) mutations.
- In combination with docetaxel, for treatment of metastatic NSCLC with disease progression on or after platinum-based chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving Cyramza.
- In combination with FOLFIRI (irinotecan, folinic acid, and 5-fluorouracil), for the treatment of metastatic colorectal cancer (CRC) with disease progression on or after prior therapy with bevacizumab, oxaliplatin, and a fluoropyrimidine.
- As a single agent, for the treatment of hepatocellular carcinoma (HCC) in patients who have an alpha fetoprotein (AFP) of ≥ 400 ng/mL and have been treated with sorafenib.

Policy/Criteria

Provider must submit documentation (including such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Cyramza is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Esophageal, Esophagogastric Junction, and Gastric Cancer (must meet all):

1. Diagnosis of esophageal, EGJ or gastric cancer;
2. Disease is unresectable, locally advanced, recurrent, or metastatic;
3. Prescribed by or in consultation with an oncologist;

4. Age \geq 18 years;
5. Prescribed as subsequent therapy in one of the following ways (a, b, or c)*:
 - a. As a single agent;
 - b. In combination with paclitaxel;
 - c. In combination with irinotecan with or without fluorouracil;**Prior authorization may be required for paclitaxel, fluorouracil or irinotecan.*
6. Request meets one of the following (a or b):*
 - a. Dose does not exceed 8 mg per kg every 2 weeks;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

B. Non-Small Cell Lung Cancer (must meet all):

1. Diagnosis of metastatic, recurrent, or advanced NSCLC;
2. Prescribed by or in consultation with an oncologist;
3. Age \geq 18 years;
4. Request meets one of the following (a or b):*
 - a. Prescribed as subsequent therapy in combination with docetaxel;
 - b. Prescribed in combination with erlotinib (Tarceva®);**Prior authorization may be required for docetaxel or erlotinib*
5. If prescribed in combination with erlotinib, disease is positive for a sensitizing EGFR mutation (e.g., EGFR exon 19 deletions or exon 21 [L858R] substitution mutation);
6. Request meets one of the following (a, b, or c):*
 - a. In combination with docetaxel: dose does not exceed 10 mg/kg on day 1 of a 21-day cycle;
 - b. In combination with erlotinib: dose does not exceed 10 mg/kg on day 1 every 2 weeks;
 - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

C. Colorectal Cancer (must meet all):

1. Diagnosis of advanced or metastatic CRC;
2. Prescribed by or in consultation with an oncologist;
3. Age \geq 18 years;
4. Prescribed in combination with irinotecan or FOLFIRI (irinotecan, folinic acid, and 5-fluorouracil);*
**Prior authorization may be required for irinotecan or FOLFIRI.*
5. Request meets one of the following (a or b):*
 - a. Dose does not exceed 8 mg/kg every 2 weeks;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

D. Hepatocellular Carcinoma (must meet all):

1. Diagnosis of HCC;
2. Prescribed by or in consultation with an oncologist;
3. Age \geq 18 years;
4. AFP \geq 400 ng/mL;
5. Disease has progressed on or after therapy with sorafenib;*
**Prior authorization may be required for sorafenib*
6. Prescribed as single-agent therapy;
7. Confirmation of Child-Pugh class A status;
8. Request meets one of the following (a or b):*
 - a. Dose does not exceed 8 mg/kg every 2 weeks;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

E. Mesothelioma (off-label) (must meet all):

1. Diagnosis of mesothelioma classified as one of the following:
 - a. Pleural;
 - b. Pericardial;
 - c. Tunica vaginalis testis;
2. Prescribed by or in consultation with an oncologist;
3. Age \geq 18 years;
4. Prescribed in combination with gemcitabine;*
**Gemcitabine may require prior authorization*
5. Prescribed as subsequent therapy;
6. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).*
**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

F. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Cyramza for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a, b, c, or d)*:

- a. Esophageal/EGJ/gastric cancer, CRC, HCC: new dose does not exceed 8 mg/kg every 2 weeks;
- b. NSCLC in combination with docetaxel: new dose does not exceed 10 mg/kg on day 1 of a 21-day cycle;
- c. NSCLC in combination with erlotinib: new dose does not exceed 10 mg/kg every 2 weeks;
- d. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy LA.PMN.53

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

AFP: alpha fetoprotein	FOLFIRI: fluorouracil, leucovorin, irinotecan
CRC: colorectal carcinoma	NCCN: National Comprehensive Cancer Network
EGJ: esophagogastric junction	NSCLC: non-small cell lung cancer
EGFR: epidermal growth factor receptor	VEGFR2: vascular endothelial growth factor receptor 2
FDA: Food and Drug Administration	
HCC: hepatocellular carcinoma	

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug	Dosing Regimen	Dose Limit/ Maximum Dose
docetaxel, paclitaxel, irinotecan, 5-FU	Esophageal, EGF, or gastric cancer: Varies	Varies
docetaxel (Taxotere®)	NSCLC: Varies	Varies
erlotinib (Tarceva®)	NSCLC: 150 mg PO QD	150 mg/day
irinotecan (Camptosar®), FOLFIRI (5-FU, leucovorin, irinotecan),	CRC: Varies	Varies

Drug	Dosing Regimen	Dose Limit/ Maximum Dose
FOLFOX (5-FU, leucovorin, oxaliplatin), CAPEOX (capecitabine, oxaliplatin)		
sorafenib (Nexavar®)	HCC: 400 mg PO BID	800 mg/day
cisplatin + pemetrexed ± bevacizumab, Opdivo® (nivolumab)/Yervoy® (ipilimumab), cisplatin + gemcitabine	Mesothelioma: Varies	Varies

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

None reported

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Gastric or EGJ adenocarcinoma	8 mg/kg IV every 2 weeks as a single agent or in combination with weekly paclitaxel	8 mg/kg
NSCLC	10 mg/kg IV on day 1 of a 21-day cycle prior to docetaxel 10 mg/kg IV every 2 weeks with daily erlotinib	10 mg/kg
CRC	8 mg/kg IV every 2 weeks prior to FOLFIRI	8 mg/kg
HCC	8 mg/kg IV every 2 weeks	8 mg/kg

VI. Product Availability

Single-dose vial: 100 mg/10 mL (10 mg/mL) solution, 500mg/50mL (10mg/mL) solution

VII. References

1. Cyramza Prescribing Information. Indianapolis, IN: Eli Lilly and Company; March 2022. Available at <http://uspl.lilly.com/cyramza/cyramza.html>. Accessed September 26, 2023.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at nccn.org. Accessed November 5, 2023.
3. National Comprehensive Cancer Network Guidelines. Esophageal and Esophagogastric Junction Cancers Version 3.2023. Available at https://www.nccn.org/professionals/physician_gls/pdf/esophageal.pdf. Accessed November 5, 2023.
4. National Comprehensive Cancer Network Guidelines. Gastric Cancer Version 2.2023. Available at https://www.nccn.org/professionals/physician_gls/pdf/gastric.pdf. Accessed November 5, 2023.
5. National Comprehensive Cancer Network Guidelines. Non-Small Cell Lung Cancer Version 4.2023. Available at https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf. Accessed November 5, 2023.

6. National Comprehensive Cancer Network Guidelines. Colon Cancer Version 3.2023. Available at https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf. Accessed November 5, 2023.
7. National Comprehensive Cancer Network Guidelines. Hepatocellular carcinoma Version 2.2023. Available at https://www.nccn.org/professionals/physician_gls/pdf/hcc.pdf. Accessed November 5, 2023.
8. National Comprehensive Cancer Network Guidelines. Mesothelioma: Pleural Version 1.2023. Available at https://www.nccn.org/professionals/physician_gls/pdf/meso_pleural.pdf. Accessed November 5, 2023.
9. Zhu AX, Kang YK, Yen CJ, et al. Ramucirumab after sorafenib in patients with advanced hepatocellular carcinoma and increased alpha-fetoprotein concentrations (REACH-2): a randomized, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2019; 20:282-96.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9308	Injection, ramucirumab, 5mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy.	01.21	04.28.21
Revised criteria for advanced esophageal, EGJ or gastric cancer allowing combination with irinotecan with or without fluorouracil and added requirement for unresectable, locally advanced, recurrent, or metastatic disease per NCCN; updated Appendix B Therapeutic Alternatives; references reviewed and updated.	09.22	10.30.22
Template changes applied to other diagnoses/indications. For esophageal, EGJ and gastric cancers, removed the requirement for “advanced” to limit possible confusion as specific disease qualifiers are outlined in the next criterion; Per NCCN Compendium, added requirements for confirmation of Child-Pugh class A status for HCC and use as single-agent therapy; for HCC, removed “progressive” cancer requirement as there is already a requirement for progression on or after sorafenib; updated Appendix B therapies; references reviewed and updated.	06.09.23	10.05.23
Added verbiage this policy is for medical benefit only.		
Annual review; per NCCN added off-label indication criteria for mesothelioma; references reviewed and updated.	05.02.24	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademarks exclusively owned by Louisiana Healthcare Connections.