

## Clinical Policy: Brentuximab Vedotin (Adcetris)

Reference Number: LA.PHAR.303

Effective Date: 07.23.22

Last Review Date: 05.13.24

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

Brentuximab vedotin for injection (Adcetris®) is a CD30-directed antibody-drug conjugate.

**\*\*Please note: This policy is for medical benefit\*\***

### FDA Approved Indication(s)

Adcetris is indicated for the treatment of adult patients with:

- Classical Hodgkin lymphoma:
  - Previously untreated Stage III or IV classical Hodgkin lymphoma (cHL), in combination with doxorubicin, vinblastine, and dacarbazine
  - cHL at high risk of relapse or progression as post-autologous hematopoietic stem cell transplantation (auto-HSCT) consolidation
  - cHL after failure of auto-HSCT or after failure of at least two prior multi-agent chemotherapy regimens in patients who are not auto-HSCT candidates
- T-cell lymphomas:
  - Previously untreated systemic anaplastic large cell lymphoma (sALCL) or other CD30-expressing peripheral T-cell lymphomas (PTCL), including angioimmunoblastic T-cell lymphoma and PTCL not otherwise specified, in combination with cyclophosphamide, doxorubicin, and prednisone
  - sALCL after failure of at least one prior multiagent chemotherapy regimen
- Primary cutaneous lymphomas:
  - Primary cutaneous anaplastic large cell lymphoma (pcALCL) or CD30-expressing mycosis fungoides (MF) who have received prior systemic therapy

Adcetris is indicated for the treatment of pediatric patients 2 years old and older with:

- Classical Hodgkin lymphoma:
  - Previously untreated high risk cHL, in combination with doxorubicin, vincristine, etoposide, prednisone, and cyclophosphamide

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of Louisiana HealthCare Connections® that Adcetris is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Classical Hodgkin Lymphoma in Adults (must meet all):

## CLINICAL POLICY

### Brentuximab Vedotin

1. Diagnosis of cHL;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age  $\geq$  18 years\*;
  - \*If age is between 2 to 21 years, consider using I.B cHL in Pediatric and Adolescent Patients below.
4. If previously untreated disease, prescribed in one of the following ways (a or b):
  - a. In combination with AVD (doxorubicin, vinblastine, and dacarbazine);
  - b. For age  $>$  60 years: In combination with dacarbazine;
5. If relapsed or refractory disease, prescribed in one of the following ways (a-e):
  - a. As a single agent;
  - b. In combination with bendamustine;
  - c. In combination with ICE (ifosfamide, carboplatin, etoposide);
  - d. In combination with nivolumab;
  - e. Following high-dose therapy and autologous stem cell rescue;
6. Request meets one of the following (a or b):\*\*
  - a. Dose does not exceed (i, ii, or iii):
    - i. Previously untreated Stage III or IV cHL: 1.2 mg/kg up to 120 mg every 2 weeks for a maximum of 12 doses;
    - ii. cHL consolidation: 1.8 mg/kg up to 180 mg every 3 weeks for a maximum of 16 cycles;
    - iii. Relapsed cHL: 1.8 mg/kg up to 180 mg every 3 weeks;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

\*\*Prescribed regimen must be FDA-approved or recommended by NCCN

#### **Approval duration: 6 months**

#### **B. Classical Hodgkin Lymphoma in Pediatric and Adolescent Patients (must meet all):**

1. Diagnosis of cHL;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age between 2 years to 21 years;
4. One of the following (a, b, or c):
  - a. If previously untreated: Prescribed as a component of Bv-AVE-PC (brentuximab vedotin, doxorubicin, vincristine, etoposide, prednisone, cyclophosphamide) or AEPA (brentuximab vedotin, etoposide, prednisone, doxorubicin);
  - b. If following AEPA: Prescribed as a component of CAPDAC (cyclophosphamide, brentuximab vedotin, prednisone, dacarbazine);
  - c. For relapsed or refractory disease (i or ii):
    - i. Prescribed in combination with involved-site radiation therapy (ISRT) or bendamustine/nivolumab/gemcitabine;
    - ii. Prescribed following high-dose therapy and autologous stem cell rescue;
5. For all requests except when prescribed in combination with ISRT or bendamustine/nivolumab/gemcitabine: Disease is classified as high risk (*see Appendix D*);
6. Request meets one of the following (a or b):\*
  - a. Dose does not exceed: 1.8 mg/kg up to 180 mg every 3 weeks for a maximum of 5 doses;

## CLINICAL POLICY

### Brentuximab Vedotin

- b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

#### **Approval duration: 6 months**

### **C. T-Cell Lymphomas (must meet all):**

1. Diagnosis of one of the following (a, b, c, d, or e):
  - a. PTCL - any of the following subtypes/histologies (i or ii):
    - i. sALCL;
    - ii. PTCL, including but not limited to the following (1, 2, 3, 4, or 5):
      - 1) Angioimmunoblastic T-cell lymphoma;
      - 2) Enteropathy-associated T-cell lymphoma;
      - 3) Monomorphic epitheliotropic intestinal T-cell lymphoma;
      - 4) Nodal PTCL with TFH phenotype;
      - 5) Follicular T-cell lymphoma;
  - b. Breast implant-associated ALCL (off-label);
  - c. Adult T-cell leukemia/lymphoma (off-label);
  - d. Relapsed or refractory extranodal NK/T-cell lymphoma (off-label);
  - e. Hepatosplenic T-cell lymphoma after failure of two first-line therapy regimens (off-label);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age  $\geq$  18 years;
4. For all requests except ALCL: Disease is CD30-positive;
5. Prescribed as a single agent or in combination with CHP (cyclophosphamide, doxorubicin, prednisone);
6. Request meets one of the following (a, b, or c):\*
  - a. Previously untreated sALCL or other CD30-positive PTCL including angioimmunoblastic T-cell lymphoma: Dose does not exceed 1.8 mg/kg up to 180 mg every 3 weeks with each cycle of chemotherapy for 6 to 8 doses;
  - b. Relapsed sALCL: Dose does not exceed 1.8 mg/kg up to 180 mg every 3 weeks;
  - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

#### **Approval duration: 6 months**

### **D. Primary Cutaneous CD30+ T-Cell Lymphoproliferative Disorders (must meet all):**

1. Diagnosis of one of the following (a, b, or c):
  - a. pcALCL;
  - b. Cutaneous ALCL with multifocal lesions or lymph node positive (off-label);
  - c. Lymphomatoid papulosis - as subsequent therapy for relapsed/refractory disease (off-label);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age  $\geq$  18 years;
4. Disease is CD30-positive;
5. Request meets one of the following (a or b):\*

## CLINICAL POLICY

### Brentuximab Vedotin

- a. Relapsed pcALCL: Dose does not exceed 1.8 mg/kg up to 180 mg every 3 weeks for a maximum of 16 cycles;
- b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

#### **E. Mycosis Fungoides/Sezary Syndrome (must meet all):**

1. Diagnosis of MF or Sezary syndrome (off-label);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age  $\geq$  18 years;
4. Request meets one of the following (a or b):\*
  - a. Relapsed CD30-positive MF: Dose does not exceed 1.8 mg/kg up to 180 mg every 3 weeks for a maximum of 16 cycles;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

#### **F. B-Cell Lymphomas (off-label) (must meet all):**

1. Diagnosis of one of the following (a, b, c, or d):
  - a. Diffuse large B-cell lymphoma;
  - b. High-grade B-cell lymphoma;
  - c. HIV-related B-cell lymphoma;
  - d. Monomorphic post-transplant lymphoproliferative disorder (PTLD) (B- or T-cell type);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. One of the following (a or b):
  - a. Age  $\geq$  18 years;
  - b. Age  $<$  18 years and both of the following (i and ii):
    - i. Relapsed or refractory primary mediastinal large B-cell lymphoma;
    - ii. Prescribed in combination with nivolumab or pembrolizumab;
4. Disease is CD30-positive;
5. For subtypes other than monomorphic PTLD (T-cell type), both of the following (a and b):
  - a. Adcetris is prescribed as subsequent therapy;
  - b. Member is not a candidate for allogeneic or autologous stem cell transplant;
6. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).\*

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

#### **G. Other diagnoses/indications (must meet 1 or 2):**

## CLINICAL POLICY

### Brentuximab Vedotin

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53

## II. Continued Therapy

### A. All Indications in Section I (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connection benefit, or documentation supports that member is currently receiving Adcetris for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):\*
  - a. New dose does not exceed (i, ii, iii, iv, v, vi, vii, or viii):
    - i. Previously untreated Stage III or IV cHL in adults: 1.2 mg/kg up to 120 mg every 2 weeks for a maximum of 12 doses;
    - ii. Previously untreated high risk cHL in pediatric and adolescent patients: 1.8 mg/kg up to 180 mg every 3 weeks for a maximum of 5 doses;
    - iii. cHL consolidation in adults: 1.8 mg/kg up to 180 mg every 3 weeks for a maximum of 16 cycles;
    - iv. Relapsed cHL in adults: 1.8 mg/kg up to 180 mg every 3 weeks;
    - v. Previously untreated sALCL or other CD30-positive PTCL including angioimmunoblastic T-cell lymphoma in adults: 1.8 mg/kg up to 180 mg every 3 weeks with each cycle of chemotherapy for 6 to 8 doses;
    - vi. Relapsed sALCL in adults: 1.8 mg/kg up to 180 mg every 3 weeks;
    - vii. Relapsed pcALCL in adults: 1.8 mg/kg up to 180 mg every 3 weeks for a maximum of 16 cycles;
    - viii. Relapsed CD30-positive MF in adults: 1.8 mg/kg up to 180 mg every 3 weeks for a maximum of 16 cycles;
  - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

\*Prescribed regimen must be FDA-approved or recommended by NCCN

### Approval duration: 12 months

### B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255

If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 2 above does not apply, refer to the off-label use policy LA.PMN.53

## III. Diagnoses/Indications for which coverage is NOT authorized

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53 for Medicaid or evidence of coverage documents.

## CLINICAL POLICY

### Brentuximab Vedotin

#### IV. Appendices/General Information

##### *Appendix A: Abbreviation/Acronym Key*

cHL: classical Hodgkin lymphoma

FDA: Food and Drug Administration

HSCT: hematopoietic stem cell transplantation

ISRT: involved-site radiation therapy

MF: mycosis fungoides

NCCN: National Comprehensive Cancer Network

pcALCL: primary cutaneous anaplastic large cell lymphoma

PTCL: peripheral T-cell lymphoma

sALCL: systemic anaplastic large cell lymphoma

SS: Sezary syndrome

##### *Appendix B: Therapeutic Alternatives*

Not applicable

##### *Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): concomitant use with bleomycin due to pulmonary toxicity
- Boxed warning(s): progressive multifocal leukoencephalopathy

##### *Appendix D: Definitions of High Risk Disease*

Per NCCN, high risk disease is defined as:

- Stage IIB with bulk disease\*

*\*Large mediastinal adenopathy (LMA): a mediastinal mass where the tumor diameter is > 1/3 the maximal thoracic diameter on an upright posteroanterior (PA) chest radiograph OR large extra-mediastinal nodal aggregate: a contiguous extramediastinal nodal aggregate that measures > 6 cm in the longest transverse diameter (transaxial measurement) or craniocaudal dimension (measured on reformatted computed tomography)*

- Stage IIIA

- Stage IIIB with E-lesions\*\*

*\*\*Localized involvement of extralymphatic tissue (by contiguous growth from an involved lymph node or in close anatomic relation) that is treatable by irradiation*

- Stage IV

Per the Adcetris pediatric cHL pivotal study, high risk was defined as the following Ann Arbor stages:

- Stage IIB with bulk disease (*see definition of bulk disease above*)
- Stage IIIB
- Stage IVA
- Stage IVB

#### V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Previously untreated Stage III	1.2 mg/kg IV up to a maximum of 120 mg in combination with chemotherapy. Administer every 2	120 mg every 2 weeks up to 12 doses

## CLINICAL POLICY

### Brentuximab Vedotin

Indication	Dosing Regimen	Maximum Dose
or IV cHL in adults	weeks until a maximum of 12 doses, disease progression, or unacceptable toxicity.	
Previously untreated high risk cHL in pediatric and adolescent patients	1.8 mg/kg IV up to a maximum of 180 mg in combination with chemotherapy. Administer every 3 weeks with each cycle of chemotherapy for a maximum of 5 doses, disease progression, or unacceptable toxicity.	180 mg every 3 weeks up to 5 doses
cHL consolidation in adults	1.8 mg/kg IV up to a maximum of 180 mg. Initiate Adcetris treatment within 4-6 weeks post-autoHSCT or upon recovery from auto-HSCT. Administer every 3 weeks until a maximum of 16 cycles, disease progression, or unacceptable toxicity.	180 mg every 3 weeks up to 16 cycles
Relapsed cHL in adults	1.8 mg/kg IV up to a maximum of 180 mg. Administer every 3 weeks until disease progression or unacceptable toxicity.	180 mg every 3 weeks
Previously untreated sALCL or other CD30-expressing PTCLs in adults	1.8 mg/kg IV up to a maximum of 180 mg in combination with cyclophosphamide, doxorubicin, and prednisone. Administer every 3 weeks with each cycle of chemotherapy for 6 to 8 doses.	180 mg every 3 weeks up to 6 to 8 doses
Relapsed sALCL in adults	1.8 mg/kg IV up to a maximum of 180 mg. Administer every 3 weeks until disease progression or unacceptable toxicity.	180 mg every 3 weeks
Relapsed pcALCL or CD30-expressing MF in adults	1.8 mg/kg IV up to a maximum of 180 mg. Administer every 3 weeks until a maximum of 16 cycles, disease progression, or unacceptable toxicity.	180 mg every 3 weeks up to 16 cycles

#### VI. Product Availability

Single-use vial: 50 mg for reconstitution

#### VII. References

1. Adcetris Prescribing Information. Bothell, WA: Seagen, Inc.; November 2022. Available at: <http://adcetrisupdate.com/>. Accessed May 17, 2023.
2. Castellino SM, et al. Brentuximab vedotin with chemotherapy in pediatric high-risk Hodgkin's lymphoma. *New Engl J Med* 2022; 387(18):1649-1660.
3. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at [www.nccn.org](http://www.nccn.org). Accessed May 17, 2023.
4. National Comprehensive Cancer Network. Hodgkin Lymphoma Version 2.2023. Available at [https://www.nccn.org/professionals/physician\\_gls/pdf/hodgkins.pdf](https://www.nccn.org/professionals/physician_gls/pdf/hodgkins.pdf). Accessed May 17, 2023.



## CLINICAL POLICY

### Brentuximab Vedotin

5. National Comprehensive Cancer Network. Pediatric Hodgkin Lymphoma Version 2.2023. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/ped\\_hodgkin.pdf](https://www.nccn.org/professionals/physician_gls/pdf/ped_hodgkin.pdf). Accessed May 17, 2023.
6. National Comprehensive Cancer Network. Primary Cutaneous Lymphomas Version 1.2023. Available at [https://www.nccn.org/professionals/physician\\_gls/pdf/primary\\_cutaneous.pdf](https://www.nccn.org/professionals/physician_gls/pdf/primary_cutaneous.pdf). Accessed May 17, 2023.
7. National Comprehensive Cancer Network. T-Cell Lymphomas Version 1.2023. Available at [https://www.nccn.org/professionals/physician\\_gls/pdf/t-cell.pdf](https://www.nccn.org/professionals/physician_gls/pdf/t-cell.pdf). Accessed May 17, 2023.
8. National Comprehensive Cancer Network. B-Cell Lymphomas Version 3.2023. Available at [https://www.nccn.org/professionals/physician\\_gls/pdf/b-cell.pdf](https://www.nccn.org/professionals/physician_gls/pdf/b-cell.pdf). Accessed May 17, 2023.

### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9042	Injection, brentuximab vedotin, 1 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy	04.22	07.23.22
Per NCCN Compendium clarified extranodal NK/T-cell lymphoma should be in the relapsed or refractory setting and removed requirement for nasal type; clarified hepatosplenic T-cell lymphoma should be after two first-line therapy regimens; references reviewed and updated.	06.02.23	10.05.23
New indication of previously untreated high risk cHL in pediatric and adolescent patients added to policy. Template changes applied to other diagnoses/indications and continued therapy section.		
Annual review: for adult cHL, added specific regimens for use per both FDA and NCCN; for pediatric cHL, moved specific staging requirements for high risk disease to Appendix D to also allow for NCCN high risk definition and updated criteria per NCCN, including requirements for use in combination with chemotherapy as well as allowance for use as subsequent therapy; for T-cell lymphomas, clarified that CD30-positive disease requirement does not apply to ALCL and added requirement for use as a single agent or in combination with CHP per NCCN; for cutaneous ALCL, added pathway for disease multifocal lesions per NCCN; for MF/SS, removed requirement for CD30-positive disease per NCCN; for B-	05.13.24	



## CLINICAL POLICY

### Brentuximab Vedotin

Reviews, Revisions, and Approvals	Date	LDH Approval Date
cell lymphomas, removed specific subtypes of DLBCL to simplify criteria, revised “AIDS-related” to “HIV-related”, added B-cell type monomorphic PTLT, added pathway for pediatric primary mediastinal large B-cell lymphoma, and added that member is not a transplant candidate for all requests except T-cell type monomorphic PTLT per NCCN; references reviewed and updated.		

#### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

## CLINICAL POLICY

### Brentuximab Vedotin

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2024 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.