

Clinical Policy: ADAMTS13, Recombinant-krhn (Adzynma)

Reference Number: LA.PHAR.635

Effective Date: 08.14.24 Last Review Date: 01.07.25 Line of Business: Medicaid

Coding Implications
Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Please note: This policy is for medical benefit

Description

ADAMTS13, recombinant-krhn (Adzynma®) is a human recombinant "A disintegrin and metalloproteinase with thrombospondin motifs 13" (rADAMTS13).

FDA Approved Indication(s)

Adzynma is indicated for prophylactic or on demand enzyme replacement therapy in adult and pediatric patients with congenital thrombotic thrombocytopenic purpura (cTTP).

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Adzynma is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Congenital Thrombotic Thrombocytopenic Purpura (must meet all):

- 1. Diagnosis of severe cTTP confirmed by all of the following (a, b, and c):
 - a. Genetic test confirming biallelic ADAMTS13 mutation;
 - b. ADAMTS13 activity < 10 % of normal, unless member is currently receiving prophylactic plasma therapy;
 - c. One of the following (i or ii):
 - i. Absence of ADAMTS13 functional inhibitor;
 - ii. Absence of anti-ADAMTS13 antibodies;
- 2. Prescribed by or in consultation with a hematologist;
- 3. Age \geq 2 years;
- 4. Failure of plasma therapy (i.e., plasma infusion, therapeutic plasma exchange), unless contraindicated or clinically significant adverse effects are experienced (*see Appendix D*);
- 5. For prophylaxis requests, member has TTP signs or symptoms that are persistent or recurrent (*see Appendix D*);
- 6. For acute (on demand) treatment requests, member has an acute TTP event defined by both the following (a and b):



- a. Platelet count $< 100,000/\mu L$ or a drop in platelet count $\ge 50\%$ of the baseline platelet count;
- b. Microangiopathic hemolytic anemia with a lactate dehydrogenase (LDH) elevation greater than two times the baseline or two times the upper limit of normal:
- 7. Documentation of member's current body weight (in kg);
- 8. Dose does not exceed the following (a or b):
 - a. For prophylactic therapy: 40 IU/kg once weekly;
 - b. For acute treatment: 40 IU/kg on day 1, followed by 20 IU/kg on day 2, and 15 IU/kg/day until two days after the acute event is resolved.

Approval duration: Prophylaxis: 6 months

Acute treatment: Up to 2 weeks per acute episode

B. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - For drugs on the PDL, the no coverage criteria policy: LA.PMN.255 for Medicaid; or
 - b. For drugs NOT on the PDL, the non-formulary policy: LA.PMN.16 for Medicaid; or
- If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy: LA.PMN.53 for Medicaid.

II. Continued Therapy

A. Congenital Thrombotic Thrombocytopenic Purpura (must meet all):

- 1. Member is currently receiving medication via Louisiana Healthcare Connections benefit or member has previously met initial approval criteria;
- 2. Member is responding positively to therapy as evidenced by, including but not limited to, improvement in <u>any</u> of the following parameters (a-c):
 - a. Thrombocytopenia;
 - b. Microangiopathic hemolytic anemia;
 - c. Symptom improvement (e.g., less headaches, lethargy, and/or abdominal pain);
- 3. Documentation of member's current body weight (in kg);
- 4. If request is for a dose increase, new dose does not exceed the following (a or b):
 - a. For prophylactic therapy: 40 IU/kg once weekly;
 - b. For acute treatment, both (i and ii):
 - i. 40 IU/kg on day 1, followed by 20 IU/kg on day 2, and 15 IU/kg/day until two days after the acute event is resolved;
 - ii. If request exceeds 2 weeks of treatment: provider justification for continued acute dosing.

Approval duration:

Prophylaxis: 12 months



Acute treatment: Up to 2 weeks per acute episode

B. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the PDL, the no coverage criteria policy: LA.PMN.255 for Medicaid; or
 - b. For drugs NOT on the PDL, the non-formulary policy: LA.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy: LA.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- **A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy LA.PMN.53 for Medicaid or evidence of coverage documents;
- **B.** Acquired/immune thrombotic thrombocytopenic purpura.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

cTTP: congenital thrombotic thrombocytopenic purpura

FDA: Food and Drug Administration

LDH: lactate dehydrogenase

rADAMTS13: a disintegrin and

metalloproteinase with thrombospondin

motifs 13

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
 Plasma infusion Fresh frozen plasma Solvent/detergent plasma Thawed plasma Plasma frozen within 24 hours of collection 	10-15 mL/kg at a frequency of every 1-3 weeks for maintenance therapy or daily for a symptomatic patient until the symptoms resolve and normalization of platelet counts	Varies

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

• Contraindication(s): life threatening hypersensitivity reactions to Adzynma or its components



• Boxed warning(s): none

Appendix D: General Information

- Examples of failure with plasma therapy in cTTP include, but are not limited to, previous stroke, kidney failure, persistent thrombocytopenia, recurrent microangiopathic hemolytic anemia, and persistent neonatal hyperbilirubinemia.
 - o Microangiopathic hemolytic anemia is a descriptive term for non-immune hemolytic anemia from intravascular red blood cell fragmentation.
- Examples of TTP signs and symptoms include, but are not limited to, persistent thrombocytopenia, recurrent microangiopathic hemolytic anemia, proteinuria, stroke, transient ischemic attack, lethargy, headaches, loss of concentration, and abdominal discomfort.
- Treatment for an acute TTP episode depends on the duration of an episode. Per the
 International Hereditary Thrombotic Thrombocytopenic Purpura Registry (van Dorland
 et al, 2019), the median duration of an acute episode was seven days. Per the Adzynma
 pivotal trial, the duration of an acute episode for Adzynma treatment ranged from 2-4
 days.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
cTTP	Prophylactic therapy	Prophylactic
	40 IU/kg IV every 2 weeks. Dosing frequency may	therapy: 40
	be adjusted to 40 IU/kg body weight once weekly	IU/kg/week
	based on prior prophylactic dosing regimen or	
	clinical response.	On-demand
		therapy: 40
	On-demand therapy	IU/kg/day
	40 IU/kg IV on day 1, followed by 20 IU/kg IV on	
	day 2, and then 15 IU/kg/day on day 3 and beyond	
	until two days after the acute event is resolved. Dose	
	is based on body weight.	

VI. Product Availability

Lyophilized powder in single-dose vials: 500 IU, 1,500 IU

VII. References

- 1. Adzynma Prescribing Information. Lexington, MA: Takeda Pharmaceuticals USA, Inc.; November 2023. Available at:
 - https://content.takeda.com/?contenttype=PI&product=ADZ&language=ENG&country=USA &documentnumber=1. Accessed December 1, 2023.
- 2. ClinicalTrials.gov. A study of BAX 930 in children, teenagers, and adults born with thrombotic thrombocytopenic purpura (TTP). Last updated February 28, 2023. Available at: https://clinicaltrials.gov/ct2/show/NCT03393975. Accessed December 1, 2023.
- 3. Alwan F, Vendramin C, Liesner R, et al. Characterization and treatment of congenital thrombotic thrombocytopenic purpura. Blood. 2019;133(15):1644-1651.



- 4. Scully M, Cataland S, Coppo P, et al. Consensus on the standardization of terminology in thrombotic thrombocytopenic purpura and related thrombotic microangiopathies. J Thromb Haemost. 2017;15(2):312-322.
- 5. van Dorland HA, Taleghani MM, Sakai K, et al. The International Hereditary Thrombotic Thrombocytopenic Purpura Registry: Key findings at enrollment until 2017. Haematologica. 2019;104(10):2107-2115.
- 6. Kremer Hovinga JA, George JN. Hereditary thrombotic thrombocytopenic purpura. N Engl J Med. 2019;381(17):1653-1662.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J7171	Injection, adamts13, recombinant-krhn, 10 iu

Reviews, Revisions, and Approvals	Date	LDH
		Approval Date
Converted corporate to local policy.	03.28.24	07.10.24
Added HCPCS code [J7171] and removed HCPCS code [C9167].	01.07.25	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. Louisiana Healthcare Connections makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Louisiana Healthcare Connections administrative policies and procedures.

This clinical policy is effective as of the date determined by Louisiana Healthcare Connections. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If



there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. Louisiana Healthcare Connections retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom Louisiana Healthcare Connections has no control or right of control. Providers are not agents or employees of Louisiana Healthcare Connections.

This clinical policy is the property of Louisiana Healthcare Connections. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2024 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.