

## Clinical Policy: Brexucabtagene Autoleucel (Tecartus)

Reference Number: LA.PHAR.472

Effective Date: 09.29.23

Last Review Date: 03.22.24

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

**\*\*Please note: This policy is for medical benefit\*\***

### Description

Brexucabtagene autoleucel (Tecartus<sup>®</sup>) is a CD19-directed chimeric antigen receptor (CAR) T cell therapy.

### FDA Approved Indication(s)

Tecartus is indicated for the treatment of:

- Adult patients with relapsed or refractory mantle cell lymphoma (MCL)\*
- Adult patients with relapsed or refractory B-cell precursor acute lymphoblastic lymphoma (ALL)

---

*\*This indication is approved under accelerated approval based on overall response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.*

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

All requests reviewed under this policy **require medical director review**.

It is the policy of Louisiana Healthcare Connections<sup>®</sup> that Tecartus is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Mantle Cell Lymphoma\* (must meet all):

*\*Only for initial treatment dose; subsequent doses will not be covered.*

1. Diagnosis of relapsed or refractory MCL;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age  $\geq$  18 years;
4. Recent (within the last 30 days) absolute lymphocyte count (ALC)  $\geq$  100 cells/ $\mu$ L;
5. Member has previously received 2 to 5 prior regimens that included all of the following (a, b, and c):
  - a. Anthracycline (e.g., doxorubicin) or bendamustine-containing chemotherapy;
  - b. Anti-CD20 monoclonal antibody therapy (e.g., rituximab);
  - c. Bruton tyrosine kinase (BTK) inhibitor (e.g., Imbruvica<sup>®</sup>, Calquence<sup>®</sup>, Brukinsa<sup>™</sup>);

## CLINICAL POLICY

### Brexucabtagene Autoleucel

6. Member does not have a history of or current central nervous system (CNS) disease or CNS disorders as detected by magnetic resonance imaging [MRI] (i.e., detectable cerebrospinal fluid malignant cells or brain metastases, CNS lymphoma, seizure disorder, cerebrovascular ischemia/hemorrhage, dementia, cerebellar disease, cerebral edema, posterior reversible encephalopathy syndrome, or any autoimmune disease with CNS involvement);
7. Member does not have a history of allogeneic stem cell transplantation;
8. Member has not previously received treatment with CAR T-cell immunotherapy (e.g., Abecma<sup>®</sup>, Breyanzi<sup>™</sup>, Carvykti<sup>™</sup>, Kymriah<sup>™</sup>, Yescarta<sup>™</sup>);
9. Tecartus is not prescribed concurrently with other CAR T-cell immunotherapy (e.g., Abecma, Breyanzi, Carvykti, Kymriah, Yescarta);
10. Dose does not exceed  $2 \times 10^8$  CAR-positive viable T cells.

**Approval duration: 3 months (1 dose only, with 4 doses of tocilizumab (Actemra) if requested at up to 800 mg per dose)**

#### **B. Acute Lymphoblastic Leukemia\*** (must meet all):

*\*Only for initial treatment dose; subsequent doses will not be covered.*

1. Diagnosis of B-cell precursor ALL;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age  $\geq 18$  years;
4. Recent (within the last 30 days) ALC  $\geq 100/\mu\text{L}$ ;
5. Request meets one of the following (a or b):
  - a. Member has relapsed or refractory disease defined as one of the following (i - iv):
    - i. Primary refractory disease;
    - ii. First relapse if first remission  $\leq 12$  months;
    - iii. Relapsed or refractory disease after 2 or more lines of systemic therapy;
    - iv. Relapsed following allogeneic stem cell transplantation (allo-SCT) and must be  $\geq 100$  days from allo-SCT at the time of Tecartus infusion;
  - b. Disease is relapsed or refractory, Philadelphia chromosome positive, and member has received 2 tyrosine kinase inhibitors (e.g., imatinib, Sprycel<sup>®</sup>, Tassigna<sup>®</sup>, Bosulif<sup>®</sup>, Iclusig<sup>®</sup>);

*\*Prior authorization may be required for tyrosine kinase inhibitors*

6. If previously treated with Blincyto<sup>®</sup>, documentation of CD19 tumor expression on blasts obtained from bone marrow or peripheral blood after completion of the most recent prior line of therapy;
7. Member does not have CNS-3 disease\* or have a history or presence of any CNS disorder (e.g., seizure disorder, cerebrovascular ischemia/hemorrhage, dementia, cerebellar disease, any autoimmune disease with CNS involvement, posterior reversible encephalopathy syndrome, or cerebral edema);

*\*CNS-3 disease is defined as detectable cerebrospinal blast cells in a sample of CSF with  $\geq 5$  white blood cells (WBCs) per  $\text{mm}^3$*

8. If member has CNS-2 disease\*, documentation of no clinically evident neurological changes;

*\*CNS-2 disease is defined as CSF blast cells with  $< 5$  WBC/ $\text{mm}^3$*

## CLINICAL POLICY

### Brexucabtagene Autoleucel

9. Member has not previously received treatment with CAR T-cell immunotherapy (e.g., Abecma, Breyanzi, Carvykti, Kymriah, Yescarta);
10. Tecartus is not prescribed concurrently with other CAR T-cell immunotherapy (e.g., Abecma, Breyanzi, Carvykti, Kymriah, Yescarta);
11. Dose does not exceed  $1 \times 10^8$  CAR-positive viable T cells.

**Approval duration: 3 months (1 dose only, with 4 doses of tocilizumab (Actemra) if requested at up to 800 mg per dose)**

#### C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the PDL (Medicaid), the no coverage criteria policy for the relevant line of business: LA.PMN.255 for Medicaid
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

## II. Continued Therapy

### A. All Indications in Section I

1. Continued therapy will not be authorized as Tecartus is indicated to be dosed one time only.

**Approval duration:** Not applicable

### B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

## III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – LA.PMN.53 for Medicaid, or evidence of coverage documents;
- B. MCL: History of or current CNS disease or CNS disorders as detected by MRI (i.e., detectable cerebrospinal fluid malignant cells or brain metastases, CNS lymphoma, seizure disorder, cerebrovascular ischemia/hemorrhage, dementia, cerebellar disease, cerebral edema, posterior reversible encephalopathy syndrome, or any autoimmune disease with CNS involvement);

## CLINICAL POLICY

### Brexucabtagene Autoleucel

- C. MCL: History of allo-SCT;
- D. ALL: History or presence of any CNS disorder, such as a seizure disorder, cerebrovascular ischemia/hemorrhage, dementia, cerebellar disease, any autoimmune disease with CNS involvement, posterior reversible encephalopathy syndrome, or cerebral edema;
- E. ALL: Presence of CNS-3 disease defined as detectable cerebrospinal blast cells in a sample of CSF with  $\geq 5$  white blood cells (WBCs) per  $\text{mm}^3$  with or without neurological changes.

#### IV. Appendices/General Information

##### *Appendix A: Abbreviation/Acronym Key*

ALC: absolute lymphocyte count

Allo-SCT: allogeneic stem cell transplantation

ALL: acute lymphoblastic leukemia

CAR: chimeric antigen receptor

CNS: central nervous system

CSF: cerebrospinal fluid

FDA: Food and Drug Administration

MCL: mantle cell lymphoma

MRI: magnetic resonance imaging

WBC: white blood cells

##### *Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
<b>Mantle Cell Lymphoma</b>		
HyperCVAD (cyclophosphamide, vincristine, doxorubicin, dexamethasone/methotrexate/ cytarabine) + rituximab	Varies	Varies
NORDIC (rituximab + cyclophosphamide, vincristine, doxorubicin, prednisone/rituximab + cytarabine)	Varies	Varies
RCHOP/RDHAP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone)/(rituximab, dexamethasone, cisplatin, cytarabine)	Varies	Varies
RDHA (rituximab, dexamethasone, cytarabine) + platinum (carboplatin, cisplatin, or oxaliplatin)	Varies	Varies
RCHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone)	Varies	Varies
Bendeka <sup>®</sup> (bendamustine) $\pm$ rituximab	Varies	Varies
VR-CAP (bortezomib, rituximab, cyclophosphamide, doxorubicin, prednisone)	Varies	Varies
Revlimid <sup>®</sup> (lenalidomide) + rituximab	Varies	Varies

## CLINICAL POLICY

### Brexucabtagene Autoleucel

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
bortezomib ± rituximab	Varies	Varies
lenalidomide ± rituximab	Varies	Varies
Imbruvica <sup>®</sup> (ibrutinib) ± rituximab	560 mg PO QD	560 mg/day
Calquence <sup>®</sup> (acalabrutinib)	100 mg PO BID	400 mg/day
Brukinsa <sup>®</sup> (zanubrutinib)	160 mg PO BID or 320 mg PO QD	320 mg/day
Venclexta <sup>®</sup> (venetoclax)	20 mg/day for week 1, 50 mg/day for week 2, 100 mg/day for week 3, 200 mg/day for week 4, 400 mg/day for week 5. Week 6 and thereafter: 800 mg/day	800 mg/day
<b>Acute Lymphoblastic Leukemia</b>		
imatinib mesylate (Gleevec <sup>®</sup> )	Adults with Ph+ ALL: 600 mg/day Pediatrics with Ph+ ALL: 340 mg/m <sup>2</sup> /day	Adults: 800 mg/day Pediatrics: 600 mg/day
Sprycel <sup>®</sup> (dasatinib)	Ph+ ALL: 140 mg per day	180 mg/day
Iclusig <sup>®</sup> (ponatinib)	Ph+ ALL: 45 mg per day	45 mg/day
Tasigna <sup>®</sup> (nilotinib)	Resistant or intolerant Ph+ CML-CP and CML-AP: 400 mg twice per day	800 mg/day
Bosulif <sup>®</sup> (bosutinib)	Ph+ CML: 500 mg per day	600 mg/day
Various combination regimens that may include the following: daunorubicin, doxorubicin, vincristine, dexamethasone, prednisone, pegaspargase, nelarabine, methotrexate, cyclophosphamide, cytarabine, rituximab, 6-mercaptopurine	Ph- ALL: varies	Varies

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

#### Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): none reported

## CLINICAL POLICY

### Brexucabtagene Autoleucel

- Boxed warning(s):
  - Cytokine release syndrome: do not administer Tecartus to patients with active infection or inflammatory disorders; treat severe or life-threatening cytokine release syndrome with tocilizumab or tocilizumab and corticosteroids
  - Neurologic toxicities: monitor for neurologic toxicities after treatment with Tecartus; provide supportive care and/or corticosteroids, as needed

#### *Appendix D: General Information*

- The ZUMA-2 trial included only patients with MCL and an ALC  $\geq 100$  cells/ $\mu$ L and a magnetic resonance imaging (MRI) of the brain showing no evidence of CNS lymphoma. Subjects with detectable cerebrospinal fluid malignant cells or brain metastases or with a history of CNS lymphoma were excluded. The trial also excluded patients with history or presence of CNS disorder, such as seizure disorder, cerebrovascular ischemia/hemorrhage, dementia, cerebellar disease, cerebral edema, posterior reversible encephalopathy syndrome, or any autoimmune disease with CNS involvement. Additionally, patients with a history of allogeneic stem cell transplantation or prior CAR therapy or other genetically modified T-cell therapy were excluded.
- Tecartus is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the Yescarta and Tecartus REMS Program.
- Refractory disease is defined as an inability to achieve a complete response to therapy.
- The ZUMA-3 trial in patients with ALL excluded patients with:
  - Presence of CNS-3 disease defined as detectable cerebrospinal blast cells in a sample of CSF with  $\geq 5$  WBCs per  $\text{mm}^3$  with or without neurological changes;
  - Presence of CNS-2 disease defined as detectable cerebrospinal blast cells in a sample of CSF with  $<5$  WBCs per  $\text{mm}^3$  with neurological changes;
  - History or presence of any CNS disorder, such as a seizure disorder, cerebrovascular ischemia/hemorrhage, dementia, cerebellar disease, any autoimmune disease with CNS involvement, posterior reversible encephalopathy syndrome, or cerebral edema.

#### V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
MCL	Target dose: $2 \times 10^6$ CAR-positive viable T cells per kg body weight	$2 \times 10^8$ CAR-positive viable T cells
ALL	$1 \times 10^6$ CAR-positive viable T cells/kg	$1 \times 10^8$ CAR-positive viable T cells/kg

#### VI. Product Availability

Single-dose unit infusion bag: frozen suspension of genetically modified autologous T-cells labeled for the specific recipient

## CLINICAL POLICY

### Brexucabtagene Autoleucl

#### VII. References

1. Tecartus Prescribing Information. Santa Monica, CA: Kite Pharma, Inc.; October 2021. Available at: <https://www.gilead.com/-/media/files/pdfs/medicines/oncology/tecartus/tecartus-pi.pdf>. Accessed October 6, 2023.
2. Wang M, Munoz J, Goy A, et al. KTE-X19 CAR T-Cell Therapy in Relapsed or Refractory Mantle-Cell Lymphoma. *N Engl J Med* 2020;382:1331-42.
3. National Comprehensive Cancer Network. B-cell Lymphomas Version 6.2023. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/b-cell.pdf](https://www.nccn.org/professionals/physician_gls/pdf/b-cell.pdf). Accessed October 12, 2023.
4. Shah BD, Ghobadi A, Oluwole OO, et al. KTE-X19 for relapsed or refractory adult B-cell acute lymphoblastic leukaemia: phase 2 results of the single-arm, open-label, multicentre ZUMA-3 study. *Lancet*. 2021 Jun 3; S0140-6736 (21) 01222-8.
5. National Comprehensive Cancer Network. Acute Lymphoblastic Leukemia Version 1.2024. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/all.pdf](https://www.nccn.org/professionals/physician_gls/pdf/all.pdf). Accessed October 12, 2023.

#### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
Q2053	Brexucabtagene autoleucl, up to 200 million autologous anti-cd19 car positive viable t cells, including leukapheresis and dose preparation procedures, per therapeutic dose

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Policy created	05.01.23	08.28.23
Annual review: per NCCN for Ph+ ALL, revised requirement to include relapse or refractory disease and modified verbiage from “failure of” to “member has received 2 tyrosine kinase inhibitors”; references reviewed and updated.	03.22.24	

#### Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing

## CLINICAL POLICY

### Brexucabtagene Autoleucl

this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2024 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained



**CLINICAL POLICY**  
**Brexucabtagene Autoleucel**

herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.