

Clinical Policy: Loncastuximab Tesirine-Ipyl (Zynlonta)

Reference Number: LA.PHAR.539

Effective Date: 09.29.23

Last Review Date: 09.16.24

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

****Please note: This policy is for medical benefit****

Description

Loncastuximab tesirine-ipyil (Zynlonta[®]) is a CD19-directed antibody and alkylating agent conjugate.

FDA Approved Indication(s)

Zynlonta is indicated for the treatment of adult patients with relapsed or refractory large B-cell lymphoma after two or more lines of systemic therapy, including diffuse large B-cell lymphoma (DLBCL) not otherwise specified, DLBCL arising from low-grade lymphoma, and high-grade B-cell lymphoma.

This indication is approved under accelerated approval based on overall response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections[®] that Zynlonta is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Large B-Cell Lymphoma (must meet all):

1. Diagnosis of large B-cell lymphoma (including DLBCL not otherwise specified, DLBCL arising from low-grade lymphoma, high-grade B-cell lymphoma, HIV-related DLBCL, primary effusion lymphoma, HHV8-positive DLBCL not otherwise specified, post-transplant lymphoproliferative disorders (PTLD), and histologic transformation of indolent lymphomas related to DLBCL);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age \geq 18 years;
4. Request meets one of the following (a or b):
 - a. Disease is refractory or member has relapsed after \geq 2 lines of systemic therapy (*see Appendix B*);
 - b. Member is not a candidate for transplant and request is for second-line therapy for partial response, no response, or progressive disease following

CLINICAL POLICY

Loncastuximab Tesirine-lpyl

chemoimmunotherapy in patients with histologic transformation to DLBCL (off-label);

5. Prescribed as a single agent;
6. Request meets one of the following (a or b):*
 - a. Dose does not exceed 0.15 mg/kg IV every 3 weeks for 2 cycles, then 0.075 mg/kg every 3 weeks for subsequent cycles;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

II. Continued Therapy

A. Large B-Cell Lymphoma (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Zynlonta for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 0.075 mg/kg every 3 weeks;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

CLINICAL POLICY

Loncastuximab Tesirine-lpyl

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – LA.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

DLBCL: diffuse large B-cell lymphoma

FDA: Food and Drug Administration

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Examples of First-Line Treatment Regimens		
RCHOP (Rituxan [®] (rituximab), cyclophosphamide, doxorubicin, vincristine, prednisone)	Varies	Varies
RCEPP (Rituxan [®] (rituximab), cyclophosphamide, etoposide, prednisone, procarbazine)	Varies	Varies
RCDOP (Rituxan [®] (rituximab), cyclophosphamide, liposomal doxorubicin, vincristine, prednisone)	Varies	Varies
DA-EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicine) + Rituxan [®] (rituximab)	Varies	Varies
RCEOP (Rituxan [®] (rituximab), cyclophosphamide, etoposide, vincristine, prednisone)	Varies	Varies
RGCVP (Rituxan [®] , gemcitabine, cyclophosphamide, vincristine, prednisone)	Varies	Varies
Examples of Second Line Treatment Regimens		
CEOP (cyclophosphamide, etoposide, vincristine, prednisone) ± Rituxan [®] (rituximab)	Varies	Varies
DHA (dexamethasone, cytarabine) + platinum (carboplatin, cisplatin, or oxaliplatin) ± Rituxan [®] (rituximab)		
DA-EPOCH ± Rituxan [®] (rituximab)	Varies	Varies
GDP (gemcitabine, dexamethasone, cisplatin) ± Rituxan [®] (rituximab)	Varies	Varies
gemcitabine, dexamethasone, carboplatin ± Rituxan [®] (rituximab)	Varies	Varies
GemOx (gemcitabine, oxaliplatin) ± Rituxan [®] (rituximab)	Varies	Varies
gemcitabine, vinorelbine ± Rituxan [®] (rituximab)	Varies	Varies

CLINICAL POLICY

Loncastuximab Tesirine-lpyl

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
lenalidomide ± Rituxan [®] (rituximab)	Varies	Varies
Rituxan [®] (rituximab)	Varies	Varies
ESHAP (etoposide, methylprednisolone, cytarabine, cisplatin) ± Rituxan [®] (rituximab)	Varies	Varies
ICE (ifosfamide, carboplatin, etoposide) ± Rituxan [®] (rituximab)	Varies	Varies
MINE (mesna, ifosfamide, mitoxantrone, etoposide) ± Rituxan [®] (rituximab)	Varies	Varies
Polivy [®] (polatuzumab vedotin) ± bendamustine ± Rituxan [®] (rituximab)	Varies	Varies
Monjuvi [®] (tafasitamab-cxix) + lenalidomide	Varies	Varies
Yescarta [®] (axicabtagene ciloleucel)	Varies	Varies
Breyanzi [®] (lisocabtagene maralucecl)	Varies	Varies

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

None reported

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Large B-cell lymphoma	0.15 mg/kg IV every 3 weeks for 2 cycles, then 0.075 mg/kg every 3 weeks for subsequent cycles	See regimen

VI. Product Availability

Lyophilized powder for reconstitution in a single-dose vial: 10 mg

VII. References

1. Zynlonta Prescribing Information. Murray Hill, NJ: ADC Therapeutics America; October 2022. Available at: www.zynlonta.com. Accessed May 6, 2024.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed April 26, 2024.
3. National Comprehensive Cancer Network. B-Cell Lymphomas Version 2.2024. Available at: https://www.nccn.org/professionals/physician_gls/pdf/b-cell.pdf. Accessed May 6, 2024.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CLINICAL POLICY

Loncastuximab Tesirine-lpyl

HCPCS Codes	Description
J9359	Injection, loncastuximab tesirine-lpyl, 0.075 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Policy created	05.01.23	09.28.23
Added Zynlonta prescribed as a single agent per NCCN; references reviewed and updated.	02.01.24	05.10.24
Annual review: revised language from “AIDS-related DLBCL” to “HIV-related DLBCL” to align with NCCN; added post-transplant lymphoproliferative disorders (PTLD) and histologic transformation of indolent lymphomas related to DLBCL as additional examples of large B-cell lymphoma; updated appendix B; references reviewed and updated.	09.16.24	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

CLINICAL POLICY

Loncastuximab Tesirine-lpyl

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2024 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.